



TEST PAPER OF JEE(MAIN) EXAMINATION – 2019

(Held On Saturday 12th JANUARY, 2019) TIME : 09 : 30 AM To 12 : 30 PM

CHEMISTRY






1. Iodine reacts with concentrated HNO_3 to yield Y along with other products. The oxidation state of iodine in Y, is :-

- (1) 5 (2) 3 (3) 1 (4) 7


Ans. (1)

Sol. $\text{I}_2 + 10\text{HNO}_3 \longrightarrow 2\text{HIO}_3 + 10\text{NO}_2 + 4\text{H}_2\text{O}$
In HIO_3 oxidation state of iodine is +5.

2. The major product of the following reaction is:



Ans. (3)




DIBAL-H will reduce cyanides & esters to aldehydes.

3. In a chemical reaction, $\text{A} + 2\text{B} \xrightleftharpoons{K} 2\text{C} + \text{D}$, the initial concentration of B was 1.5 times of the concentration of A, but the equilibrium concentrations of A and B were found to be equal. The equilibrium constant(K) for the aforesaid chemical reaction is :

- (1) 16 (2) 4 (3) 1 (4)
- $\frac{1}{4}$

Ans.(2)



At equilibrium $[\text{A}] = [\text{B}]$

$$a_0 - x = 1.5a_0 - 2x \Rightarrow x = 0.5a_0$$

t = t_{eq}	$0.5a_0$	$0.5a_0$	a_0	$0.5a_0$
---------------------	----------	----------	-------	----------

$$K_{\text{C}} = \frac{[\text{C}]^2 [\text{D}]}{[\text{A}] [\text{B}]^2} = \frac{(a_0)^2 (0.5a_0)}{(0.5a_0) (0.5a_0)^2} = 4$$

4. Two solids dissociate as follows





The total pressure when both the solids dissociate simultaneously is :-

(1) $x^2 + y^2 \text{ atm}$ (2) $x^2 + y^2 \text{ atm}$

(3) $2(\sqrt{x+y}) \text{ atm}$ (4) $\sqrt{x+y} \text{ atm}$

Ans. (3)



Adding (1) and (2)

$$x + y = (P_1 + P_2)^2$$

Now total pressure

$$P_{\text{T}} = P_{\text{C}} + P_{\text{B}} + P_{\text{E}}$$

$$= (P_1 + P_2) + P_1 + P_2 = 2(P_1 + P_2)$$

$$P_{\text{T}} = 2(\sqrt{x+y})$$

5. Freezing point of a 4% aqueous solution of X is equal to freezing point of 12% aqueous solution of Y. If molecular weight of X is A, then molecular weight of Y is :-

- (1) A
-
- (2) 3A
-
- (3) 4A
-
- (4) 2A

Ans. (2)



JEE (Main) Examination-2019/Morning Session/12-01-2019



Sol. For same freezing point, molality of both solution should be same.

$$m_x = m_y$$

$$\frac{4 \times 1000}{96 \times M_x} = \frac{12 \times 1000}{88 \times M_y}$$

$$\text{or, } M_y = \frac{96 \times 12}{4 \times 88} M_x = 3.27 A$$

Closest option is 3A.





6. Poly- β -hydroxybutyrate-co- β -hydroxyvalerate (PHBV) is a copolymer of ____.

- (1) 3-hydroxybutanoic acid and 4-hydroxypentanoic acid
- (2) 2-hydroxybutanoic acid and 3-hydroxypentanoic acid
- (3) 3-hydroxybutanoic acid and 2-hydroxypentanoic acid
- (4) 3-hydroxybutanoic acid and 3-hydroxypentanoic acid

Ans. (4)

Sol. PHBV is a polymer of 3-hydroxybutanoic acid and 3-Hydroxy pentanoic acid.

7. Among the following four aromatic compounds, which one will have the lowest melting point ?




Ans. (1)

Sol. M.P. of Naphthalene $\approx 80^\circ\text{C}$

8. $\text{CH}_3\text{CH}_2-\overset{\text{OH}}{\underset{\text{Ph}}{\text{C}}}-\text{CH}_3$ cannot be prepared by :

- (1) $\text{HCHO} + \text{PhCH}(\text{CH}_3)\text{CH}_2\text{MgX}$
- (2) $\text{PhCOCH}_2\text{CH}_3 + \text{CH}_3\text{MgX}$
- (3) $\text{PhCOCH}_3 + \text{CH}_3\text{CH}_2\text{MgX}$
- (4) $\text{CH}_3\text{CH}_2\text{COCH}_3 + \text{PhMgX}$

Ans. (1)



9. The volume of gas A is twice that of gas B. The compressibility factor of gas A is thrice than that of gas B at same temperature. The pressures of the gases for equal number of moles are :

- (1) $2P_A = 3P_B$
- (2) $P_A = 3P_B$
- (3) $P_A = 2P_B$
- (4) $3P_A = 2P_B$

Ans. (1)

Sol. $V_A = 2V_B$
 $Z_A = 3Z_B$

$$\frac{P_A V_A}{n_A R T_A} = \frac{3 \cdot P_B \cdot V_B}{n_B \cdot R T_B}$$

$$2P_A = 3P_B$$

10. The element with $Z = 120$ (not yet discovered) will be an/a :

- (1) transition metal
- (2) inner-transition metal
- (3) alkaline earth metal
- (4) alkali metal

Ans. (3)

Sol. $Z = 120$

Its general electronic configuration may be represented as [Nobal gas] ns^2 , like other alkaline earth metals.



JEE (Main) Examination-2019/Morning Session/12-01-2019


11. Decomposition of X exhibits a rate constant of 0.05 $\mu\text{g}/\text{year}$. How many years are required for the decomposition of 5 μg of X into 2.5 μg ?
 (1) 50 (2) 25 (3) 20 (4) 40

Ans. (1)


Sol. Rate constant (K) = 0.05 $\mu\text{g}/\text{year}$ means zero order reaction

$$t_{1/2} = \frac{a_0}{2K} = \frac{5\mu\text{g}}{2 \times 0.05\mu\text{g}/\text{year}} = 50 \text{ year}$$

12. The major product of the following reaction is :



Ans. (4)



13. Given

Gas	H ₂	CH ₄	CO ₂	SO ₂
Critical Temperature/K	33	190	304	630


On the basis of data given above, predict which of the following gases shows least adsorption on a definite amount of charcoal ?

- (1) H₂ (2) CH₄ (3) SO₂ (4) CO₂

Ans. (1)

Sol. Smaller the value of critical temperature of gas, lesser is the extent of adsorption. so least adsorbed gas is H₂

14. For diatomic ideal gas in a closed system, which of the following plots does not correctly describe the relation between various thermodynamic quantities ?



Ans. (2)

Sol. At higher temperature, rotational degree of freedom becomes active.

$$C_p = \frac{7}{2}R \quad (\text{Independent of } P)$$


$$C_v = \frac{5}{2}R \quad (\text{Independent of } V)$$

Variation of U vs T is similar as C_v vs T.

15. The standard electrode potential E[⊖] and its

temperature coefficient $\left(\frac{dE^\ominus}{dT}\right)$ for a cell are 2V

and $-5 \times 10^{-4} \text{ VK}^{-1}$ at 300 K respectively. The cell reaction is



The standard reaction enthalpy ($\Delta_r H^\ominus$) at 300

K in kJ mol⁻¹ is,

[Use R = 8J K⁻¹ mol⁻¹ and F = 96,000 C mol⁻¹]

- (1) -412.8 (2) -384.0
 (3) 206.4 (4) 192.0

Ans. (1)

Sol. Chiefly NO₂, O₃ and hydrocarbon are responsible for build up smog.

Click Here for JEE Mains & Advanced Free Online Test Practice

<https://examination-system.shineitsolutions.in/jee-mains-and-advanced-examination-system/student-login.php>



JEE (Main) Examination-2019/Morning Session/12-01-2019



16. The molecule that has minimum/no role in the formation of photochemical smog, is :

- (1) $\text{CH}_2 = \text{O}$
- (2) N_2
- (3) O_3
- (4) NO

Ans. (2)

Sol. Chiefly NO_2 , O_3 and hydrocarbon are responsible for build up smog.

17. In the Hall-Heroult process, aluminium is formed at the cathode. The cathode is made out of :

- (1) Platinum
- (2) Carbon
- (3) Pure aluminium
- (4) Copper

Ans. (2)

17. Ans.(2) Carbon

Sol. In the Hall-Heroult process the cathode is made of carbon.







18. Water samples with BOD values of 4 ppm and 18 ppm, respectively, are :

- (1) Highly polluted and Clean
- (2) Highly polluted and Highly polluted
- (3) Clean and Highly polluted
- (4) Clean and Clean

Ans. (3)

Sol. Clean water would have BOD value of less than 5 ppm whereas highly polluted water could have a BOD value of 17 ppm or more.

19. In the following reactions, products A and B are :




Ans. (4)

19.

Ans. 4

Sol.



20. What is the work function of the metal if the light of wavelength 4000 \AA generates photoelectrons of velocity $6 \times 10^5 \text{ ms}^{-1}$ from it ?

(Mass of electron = $9 \times 10^{-31} \text{ kg}$)

Velocity of light = $3 \times 10^8 \text{ ms}^{-1}$

Planck's constant = $6.626 \times 10^{-34} \text{ Js}$

Charge of electron = $1.6 \times 10^{-19} \text{ JeV}^{-1}$)

- (1) 0.9 eV
- (2) 4.0 eV
- (3) 2.1 eV
- (4) 3.1 eV

Ans. (3)



JEE (Main) Examination-2019/Morning Session/12-01-2019

Sol. $h\nu = \phi + h\nu'$

$$\frac{1}{2}mv^2 = hc\left(\frac{1}{\lambda} - \frac{1}{\lambda_0}\right)$$

$$h\nu = \phi + \frac{1}{2}mv^2$$

$$\phi = \frac{6.626 \times 10^{-34} \times 3 \times 10^8}{4000 \times 10^{-10}} - \frac{1}{2} \times 9 \times 10^{-31} \times (6 \times 10^5)^2$$

$$\phi = 3.35 \times 10^{-19} \text{ J} \Rightarrow \phi \approx 2.1 \text{ eV}$$

21. Among the following compounds most basic amino acid is :

- (1) Lysine
- (2) Asparagine
- (3) Serine
- (4) Histidine

Ans. (4)



Sol. Histidine

22. The metal d-orbitals that are directly facing the ligands in $K_3[Co(CN)_6]$ are :

- (1) d_{xz} , d_{yz} and d_{z^2}
- (2) d_{xy} , d_{xz} and d_{yz}
- (3) d_{xy} and $d_{x^2-y^2}$
- (4) $d_{x^2-y^2}$ and d_{z^2}

Ans. (4)

Sol. $K_3[Co(CN)_6]$



23. The hardness of a water sample (in terms of equivalents of $CaCO_3$) containing $10^{-3} \text{ M } CaSO_4$ is :

(molar mass of $CaSO_4 = 136 \text{ g mol}^{-1}$)

- (1) 100 ppm
- (2) 50 ppm
- (3) 10 ppm
- (4) 90 ppm

Ans. (1)

Sol. ppm of $CaCO_3$

$$(10^{-3} \times 10^3) \times 100 = 100 \text{ ppm}$$

24. The correct order for acid strength of compounds $CH \equiv CH$, $CH_3-C \equiv CH$ and $CH_2=CH_2$ is as follows :

- (1) $CH \equiv CH > CH_2 = CH_2 > CH_3-C \equiv CH$
- (2) $HC \equiv CH > CH_3-C \equiv CH > CH_2 = CH_2$
- (3) $CH_3-C \equiv CH > CH_2 = CH_2 > HC \equiv CH$
- (4) $CH_3-C \equiv CH > CH \equiv CH > CH_2 = CH_2$

Ans. (2)

Sol. $CH \equiv CH > CH_3-C \equiv CH > CH_2=CH_2$


(Acidic strength order)

25. $Mn_2(CO)_{10}$ is an organometallic compound due to the presence of :


- (1) Mn - Mn bond
- (2) Mn - C bond
- (3) Mn - O bond
- (4) C - O bond

Ans. (2)

Sol. Compounds having at least one bond between carbon and metal are known as organometallic compounds.



26. The increasing order of reactivity of the following compounds towards reaction with alkyl halides directly is :



- (1) (B) < (A) < (D) < (C)
- (2) (B) < (A) < (C) < (D)
- (3) (A) < (C) < (D) < (B)
- (4) (A) < (B) < (C) < (D)


Ans. (2)



JEE (Main) Examination-2019/Morning Session/12-01-2019




Sol. Nucleophilicity order




27. The pair of metal ions that can give a spinonly magnetic moment of 3.9 BM for the complex $[M(H_2O)_6]Cl_2$, is :

- (1) Cr^{2+} and Mn^{2+} (2) V^{2+} and Co^{2+}
 (3) V^{2+} and Fe^{2+} (4) Co^{2+} and Fe^{2+}

Ans. (2)


27. Ans.(2) V^{2+} and Co^{2+}  $3d^3$

3 unpaired e^- , spin only
 magnetic moment
 = 3.89 B.M.

 $3d^7$

3 unpaired e^- , spin only
 magnetic moment
 = 3.89 B.M.

28. In the following reaction




Aldehyde	Alcohol
HCHO	t BuOH
CH_3CHO	MeOH

The best combinations is :

- (1) HCHO and MeOH
 (2) HCHO and t BuOH
 (3) CH_3CHO and MeOH
 (4) CH_3CHO and t BuOH

Ans. (1)



$$\text{rate} \propto \frac{1}{\text{steric crowding of aldehyde}}$$

t -butanol can show formation of carbocation in acidic medium.

29. 50 mL of 0.5 M oxalic acid is needed to neutralize 25 mL of sodium hydroxide solution. The amount of NaOH in 50 mL of the given sodium hydroxide solution is :

- (1) 40 g (2) 20 g (3) 80 g (4) 10 g

BONUS



$$m_{eq} \text{ of } H_2C_2O_4 = m_{eq} \text{ NaOH}$$

$$50 \times 0.5 \times 2 = 25 \times M_{NaOH} \times 1$$

$$\therefore M_{NaOH} = 2 \text{ M}$$



$$\text{Now } 1000 \text{ ml solution} = 2 \times 40 \text{ gram NaOH}$$

$$\therefore 50 \text{ ml solution} = 4 \text{ gram NaOH}$$

30. A metal on combustion in excess air forms X, X upon hydrolysis with water yields H_2O_2 and O_2 along with another product. The metal is :

(1) Rb (2) Na (3) Mg (4) Li

Ans. (1)



Concepts Made Easy
CBSE NEET
AIIMS Exam 2019
 Free Online Exam
 Give test &
 Check your All India Rank
 www.ConceptsMadeEasy.com

Click Here for CBSE NEET &
 AIIMS Free Online Test
 Practice
<https://examination-system.shineitsolutions.in/cbse-neet-and-aiims-examination-system/student-login.php>